

SUBDUCTION INITIATION OF GEOSCIENTISTS: KECK CATALINA II

ZEB PAGE, Oberlin College

JADE STAR LACKEY, Pomona College

NICOLLETTE MITCHELL, Project Mentor, Oberlin College

CLARAMARGARET FLOOD, Peer Mentor, Oberlin College

Students: **BLAIZE ADLER-IVANBROOK**, Colorado College; **NATHANIEL BESS**, Franklin & Marshall College; **AMANI CANADA**, Trinity University; **JUAN PABLO ESPARZA**, Tarrant County College; **ERIC HASEGAWA**, Amherst College; **SARAH HAMPTON**, Oberlin College; **JAMES KARROUM II**, Rice University; **LORENA PARAS**, Smith College; **PAIGE VOSS**, Pomona College

INTRODUCTION

Franciscan-like subduction-related metamorphic rocks (blocks of garnet-bearing blueschist and amphibolite) were recognized on Santa Catalina Island by the first half of the twentieth century (Woodford, 1924; Bailey, 1941). However, the modern era of Catalina geology began with detailed mapping and a tectonic model by Platt (1975). The metamorphic rocks of Catalina (sometimes referred to as the Catalina Schist) consist of mappable units of metasedimentary and ultramafic mélange that range from lawsonite-blueschist to amphibolite facies, with increasing grade correlated with structural height (Platt, 1975). Platt's initial subdivision of the island into blueschist, greenschist, amphibolite, and ultramafic units bounded by shallow dipping thrust faults (Fig. 1) has been refined, with the "greenschist" unit now defined as epidote amphibolite and epidote blueschist overprinted with greenschist-facies assemblages (Grove and Bebout, 1995).

The highest-grade amphibolite unit from Catalina records peak conditions of 7-12 kbar and 650-750 °C based on cation thermometry and fluid inclusion barometry (Platt, 1975; Sorensen and Barton, 1987). The unusually high temperatures and Barrovian-like assemblages of the highest-grade rocks became the basis for the interpretation that the Catalina Schist was formed in a nascent subduction zone. In this model, the amphibolite-facies rocks were

formed at the initiation of subduction and recorded high temperatures due to proximity with the hot mantle wedge; the inverted metamorphic gradient of underthrust lower-grade units recorded the subsequent cooling of the trench (Platt, 1975; Cloos, 1985; Peacock, 1987). More recent analysis of detrital zircon ages from Catalina metasedimentary rocks has revealed that accretion of the Catalina Schist occurred over at least a 20 My period, with the lowest-grade units containing detrital zircons younger than the 115 Ma metamorphic age yielded by the high-grade rocks (Grove et al., 2008). Furthermore, a single garnet blueschist block (Fig. 1) found in the blueschist-facies mélange yielded high-error Rb-Sr and Ar-Ar ages of ~150 Ma suggesting that a pre-Catalina subduction zone existed in the region. The age of the garnet blueschist block was firmly established by a 155 ± 8 U-Pb sphene age collected during the 2012 Keck project (Awalt et al., 2013). However, the P-T history of this key sample remains unconstrained, possibly due to multiple episodes of subduction as the Catalina trench become superimposed on the remnants of a thermally mature subduction zone through subduction erosion (Grove et al., 2008).

Santa Catalina Island has long been used as a field laboratory to investigate devolatilization and fluid flow in subduction zones. Petrologic, geochemical, and chronological constraints on Santa Catalina Island have been used to document extensive fluid flow and




Figure 1: Geologic sketch map of Santa Catalina Island, California, modified from Platt (1975) and Grove and Barton (1995). The location of Catalina Island is shown in the red star on the inset map; the Franciscan Complex is shown in purple. The proposed study areas are shown in black rectangles.

metasomatism due to devolatilization of sediments during subduction (e.g., Sorensen and Grossman, 1989; Bebout, 1991; 2007; King et al., 2006, 2007). Many blocks in the amphibolite/ultramafic mélangé have talc or tremolite metasomatic selvages or reaction rinds similar to those found in the Franciscan Complex. Unlike in the Franciscan, some of these selvages contain garnet that is compositionally different from that found in the blocks (Penniston-Dorland et al., 2013). In situ analysis of oxygen isotopes based on samples collected during the 2012 Keck Project has demonstrated that selvages contain garnet that grew both before and after rind formation (Leung et al., 2016). These garnets likely contain further information on the P-T history of rind formation and on the trace element composition of rind forming fluids.

In the summer of 2018, we undertook a nine-student Keck Gateway project using the geology of Santa Catalina Island, California as an introduction to the

study of metamorphic rocks and metasomatism both in the laboratory and in the field. Students worked collectively on three projects using in situ analysis of major and trace elements in garnet to address questions of tectonics (timing of garnet vs. lawsonite growth in garnet blueschist, trace element zoning in amphibolite garnets with and without plagioclase) and metasomatism (major and trace elements in rind garnets). The project began with students joining Page, Buckle, and Flood on the Oberlin campus for two weeks. During this time, we had several local field trips and team-building exercises but had a primary focus on analyzing garnets from previously-prepared thick sections by SEM/EDS. By the end of the two weeks, major-element traverses of garnets from four different tectonic blocks from Catalina were completed, and targets were selected for trace-element analysis. At this point, the team flew to Los Angeles, joining Lackey for a week in the field that involved investigating field relations of the tectonic blocks already under study, limited new sample collection,

and a digital mapping exercise on blocks in mélangé. The final two weeks of the project were spent on the campus of Pomona College measuring trace-element compositions of garnet and lawsonite by laser ablation inductively-coupled mass spectrometry (LA-ICP-MS), measuring bulk-rock compositions by X-ray Fluorescence Spectrometry, analyzing data, drafting abstracts and posters.

STUDENT PROJECTS

An investigation of the Catalina garnet-blueschist: Major and trace element composition and zoning in garnet and lawsonite from multiply subducted block

Blaize Adler-Ivanbrooke, Juan Esparza, Sarah Hampton

As described above, a large garnet-lawsonite blueschist block with an anomalously old age is a key sample in understanding the tectonic history of Catalina. Preliminary attempts at thermodynamic modeling in order to reconstruct the pressure-temperature evolution of the rock did not succeed in reproducing the observed metamorphic assemblage. One possibility is that the garnet and dated sphene formed in an early metamorphism at 155Ma, and the lawsonite and glaucophane recrystallized in a second lawsonite-blueschist metamorphism after 115Ma. This is consistent with color zonation in glaucophane (Awalt et al., 2013). Detailed petrography by electron microscope revealed that lawsonite is texturally late, and, in particular, exists as veins that crosscut resorbed garnet adding further credence to a multiple-subduction origin (Adler-Ivanbrooke et al. (2018). Elevated Heavy Rare Earth Element (HREE) content in both late lawsonite and early garnet suggest that lawsonite grew at the expense of garnet or with a metasomatic source of HREE (e.g., Mulcahy et al., 2014). Garnet exists only in a m-thick band through the block, and whole rock geochemistry of multiple samples across the block show that this is not because of compositional differences and is likely due to preservation of garnet only within the core of the block.

Major and Trace Element Zoning in Garnets of Unusual Size from Blocks Hosted by Ultramafic Mélange, Santa Catalina Island, California

Amani Canada, James Karroum II, Lorena Paras

Most Catalina garnet-hornblende rocks contain no plagioclase, however, the presence of plagioclase and granitic veins in some blocks led Sorensen and Barton (1987) to conclude that all garnet-hornblende rocks are residuum from the partial melting of a basaltic precursor during subduction initiation. Canada et al. (2018) compared the major and trace-element zoning patterns in three garnet-hornblende blocks: two (one plagioclase-bearing) with typical garnet sizes for the island (<5mm diameter), and a third unusual plagioclase-bearing sample containing ~3cm diameter garnets collected during the 2012 Keck project.

Garnets from all three samples were found to have similar major and trace-element zoning patterns, suggesting a similar bulk composition and P-T evolution for mafic blocks with or without plagioclase, supporting the notion of the absence of plagioclase being due to melt extraction. Unlike the single existing published analysis of trace elements in Catalina (Sorenson and Grossman, 1989), HREE were not found to increase in the rims of garnets, however, Cr did increase in some garnet rims, perhaps due to metasomatism from a mantle-source.

Major and trace element analysis of garnet crystals from a hornblendeite block and rind on Santa Catalina Island, CA: Insights into metasomatic processes in subduction mélangé

Nathaniel Bess, Eric Hasegawa, Paige Voss

Leung et al. (2016) documented three different generations of garnets in a single rind sample from one block on Catalina, first described by Penniston-Dorland et al., (2013). Three students worked collaboratively to describe the mineralogy of a block and two samples of garnet-bearing rind (Fig. 2). Bess et al. (2018) showed that, similar to the oxygen isotope record, different populations of garnets in block and rind show different trace-element patterns. Metasomatic rinds have two different generations of garnet, mm-scale crystals that have core major and trace-element compositions similar to the block garnet. Rims on large rind garnets differ from the rims of block garnets, but are similar in major and trace elements to a second generation of 100µm-scale



Figure 2: Students draw and photograph a tectonic block of garnet hornblendite with actonite-chlorite rind at Ripper's Cove, Santa Catalina Island, California.

garnets found in quartz veins that cut the rind.

Collectively, the three groups were able to provide the community important new trace-element data on garnets to the study of metamorphism and metasomatism on Santa Catalina Island. These data help shed new light on slab-melting within nascent subduction zones, and help add more evidence for the idea of multiple episodes of subduction within the subduction mélange channel. In undertaking this research students gained valuable field and laboratory skills as well as participating in many discussions on the different careers available to geologists and the importance of and challenges inherent in increasing diversity in the geosciences.

REFERENCES

- Awalt, M.B., Page, F.Z., Walsh, E.O., Kylander-Clark, A.R.C., and Wirth, K.R., 2013, New evidence for old subduction in the Catalina Schist, Santa Catalina Island, CA: Geological Society of America Abstracts with Programs, v. 45, p. 798.
- Adler-Ivanbrook B, Hampton SK, Esparza Limon JP, Lackey JS, and Page FZ (2018) An investigation of the Catalina garnet-blueschist: Major and trace element composition and zoning in garnet and lawsonite from a multiply subducted block, Abstract T21F-0290 presented at 2018 AGU Fall Meeting, Washington, DC 10-14 Dec.
- Bailey, E.H., 1941, Mineralogy, petrology, and geology of Catalina Island, California: Ph.D. Thesis, Stanford University.
- Bebout, G.E., 1991, Field-based evidence for devolatilization in subduction zones; implications for arc magmatism: Science, v. 251, p. 413–416.
- Bebout, G.E., 2007, Metamorphic chemical geodynamics of subduction zones: Earth and Planetary Science Letters, v. 260, no. 3-4, p. 373–393.

- Bess NT, Hasegawa EM, Voss PR, Lackey JS, and Page FZ (2018) Major and trace element analysis of garnet crystals from a hornblende block and rind on Santa Catalina Island, CA: Insights into metasomatic processes in subduction mélange, Abstract T21F-0292 presented at 2018 AGU Fall Meeting, Washington, DC 10-14 Dec.
- Canada AL, Karroum JG, Paras L, Lackey JS, and Page FZ (2018) Major and Trace Element Zoning in Garnets of Unusual Size from Blocks Hosted by Ultramafic Mélange, Santa Catalina Island, California, Abstract T21F-0291 presented at 2018 AGU Fall Meeting, Washington, DC 10-14 Dec.
- Cloos, M., 1985, Thermal Evolution of Convergent Plate Margins - Thermal Modeling and Reevaluation of Isotopic Ar-Ages for Blueschists in the Franciscan Complex of California: *Tectonics*, v. 4, p. 421–433.
- Grove, M., and Bebout, G., 1995, Cretaceous tectonic evolution of coastal southern California: Insights from the Catalina Schist: *Tectonics*, v. 14, p. 1290–1308.
- Grove, M., Bebout, G.E., Jacobson, C., Barth, A., Kimbrough, D., King, R.L., Zou, H., Lovera, O., Mahoney, B., and Gehrels, G.E., 2008, The Catalina Schist: Evidence for middle Cretaceous subduction erosion of southwestern North America, in *Formation and Applications of the Sedimentary Record in Arc Collision Zones*: Geological Society of America Special Paper 436, p. 335–361.
- King, R.L., Bebout, G.E., Moriguti, T., and Nakamura, E., 2006, Elemental mixing systematics and Sr-Nd isotope geochemistry of melange formation: Obstacles to identification of fluid sources to arc volcanics: *Earth and Planetary Science Letters*, v. 246, p. 288–304.
- King, R.L., Bebout, G.E., Grove, M., Moriguti, T., and Nakamura, E., 2007, Boron and lead isotope signatures of subduction-zone melange formation: Hybridization and fractionation along the slab-mantle interface beneath volcanic arcs: *Chemical Geology*, v. 239, p. 305–322.
- Leung, M.C., Page, F.Z., Penniston-Dorland, S.C., Kitajima, K., and Valley, J.W., 2016, Constraints on garnet-bearing metasomatic rind growth in subduction mélange through SIMS analysis of oxygen isotopes, in *Geological Society of America*, p. 281880–1.
- Mulcahy, S.R., Vervoort, J., and Renne, P.R., 2014, Dating subduction-zone metamorphism with combined garnet and lawsonite Lu-Hf geochronology: *Journal of Metamorphic Geology*, v. 32, no. 5, p. 515–533, doi: 10.1111/jmg.12092.
- Peacock, S., 1987, Creation and Preservation of Subduction-Related Inverted Metamorphic Gradients: *Journal of Geophysical Research-Solid Earth and Planets*, v. 92, p. 12763–12781.
- Penniston-Dorland, S.C., Gorman, J.K., Bebout, G.E., Piccoli, P.M., and Walker, R.J., 2014, Reaction rind formation in the Catalina Schist: Deciphering a history of mechanical mixing and metasomatic alteration: *Chemical Geology*, v. 384, no. C, p. 47–61, doi: 10.1016/j.chemgeo.2014.06.024.
- Platt, J.P., 1975, Metamorphic and deformational processes in the Franciscan Complex, California: some insights from the Catalina Schist terrane: *Bulletin of the Geological Society of America*, v. 86, p. 1337–1347.
- Sorensen, S.S., 1986, Petrologic and geochemical comparison of the blueschist and greenschist units of the Catalina Schist Terrane, southern California, in Evans, B.W. and Brown, E.H. eds. *Blueschists and Eclogites*: Geological Society of America Memoir 164, Geological Society of America, p. 59–75.
- Sorensen, S.S., and Grossman, J., 1989, Enrichment of trace elements in garnet amphibolites from a paleo-subduction zone: Catalina Schist, Southern California: *Geochimica et Cosmochimica Acta*, v. 53, p. 3155–3177.
- Sorensen, S.S., and Barton, M., 1987, Metasomatism and Partial Melting in a Subduction Complex - Catalina Schist, Southern California: *Geology*, v.

15, p. 115–118.

Woodford, A.O., 1924, The Catalina metamorphic facies of the Franciscan Series: University of California Publications in Geological Sciences, v. 15, p. 49–68.